Bio& 241 A&P 1 Unit 2 / Lecture 3



Functional classification of joints

(based on degree of movement permitted within the joint)

- Synarthrosis: Immovable joints
 - *a)* Sutures = joint between bones in the skull.
 - b) Synchondrosis = epiphyseal plate.
 - (Hyaline cartilage is eventually replaced by bone).
 - c) Synostosis = epiphyseal line of mature bones.

• Amphiarthrosis: Slightly movable joints

- a) Syndesmosis = dense fibrous connective tissue located
 on the distal or proximal portion of long bones.
 (eg. distal tibia and fibula, also distal radius and ulna)
- b) Symphysis = cartilaginous joint (fibrocartilage). (eg. pubic symphysis and intervertebral discs)

Functional classification of joints

(based on degree of movement permitted within the joint)


• Diarthrosis: Freely movable joints

c) synovial joints:


have a space between the bones called a synovial joint cavity, containing synovial fluid. The joint also has articular cartilage.

Synarthrosis: immovable joints

Suture




Synarthrosis: immovable joints




Amphiarthrosis: Slightly movable joints

Symphysis



Amphiarthrosis: Slightly movable joints

Syndesmosis



Synarthrosis or Amphiarthrosis?

Gomphosis



Diarthrosis: freely movable joints



Structural classification of joints

(based on the presence or lack of a joint cavity and fibrous connective tissue)

•Fibrous Joints

NO joint cavity. Fibrous CT hold bones together.


•Cartilaginous Joints

NO joint cavity. Cartilage connects bones together

•Synovial Joints

HAVE a joint cavity, containing synovial fluid, and ligaments which provide extra support. Cartilage is also present.

Fibrous joints




Cartilaginous Joints

- Synchondroses: Epiphyseal plate

First Rib and Manubrium


- Symphyses: Pubic symphysis

Synchondrosis





O John Wiley & Sons, Inc.

Symphysis




Synovial Joints





70% of knee injuries involve stretching or tearing the ACL



Navicular
Second
cuneiform
Third
cuneiform
and third cuneiforms of the tarsus in the foot

Humerus


Trochlea


Trochlear
notch


(b) Hinge joint between trochlea of humerus
and trochlear notch of ulna at the elbow

O John Wiley & Sons, Inc.


9








(a) Atlanto-occipital and cervical intervertebral joints




(d) Wrist joint

O John Wiley & Sons, Inc.




(b) Shoulder joint

O John Wiley & Sons, Inc.



(c) Elbow joint





(e) Hip joint


(f) Knee joint

O John Wiley & Sons, Inc.




(a) Shoulder joint

O John Wiley & Sons, Inc.



(b) Wrist joint







(c) Hip joint

(d) Metacarpophalangeal joints of the fingers (not the thumb) $\,$


O John Wiley & Sons, Inc.






(a) Shoulder joint

(b) Hip joint







(a) Atlanto-axial joint


(b) Shoulder joint


O John Wiley & Sons, Inc.



(c) Hip joint







(h) Radioulnar joint